



LEADING ARTICLE

Duplex Ultrasound Investigation of the Veins of the Lower Limbs after Treatment for Varicose Veins – UIP Consensus Document **CME**

M. De Maeseneer^{a,b,*}, O. Pichot^{c,d}, A. Cavezzi^e, J. Earnshaw^f, A. van Rij^g, F. Lurie^h, P.C. Smithⁱ

^a Department of Dermatology, Erasmus Medical Centre, PB 2040, 3000 CA Rotterdam, The Netherlands

^b Surgery, Faculty of Medicine, University of Antwerp, Antwerp, Belgium

^c Department of Vascular Medicine, University Hospital Grenoble, Grenoble, France

^d Department of Vascular Surgery, University Hospital Grenoble, Grenoble, France

^e Vascular Unit, Clinica Stella Maris and Poliambulatorio Hippocrates, San Benedetto del Tronto, Marche, Italy

^f Department of Vascular Surgery, Gloucestershire Royal Hospital, Gloucester, UK

^g Department of Surgery, Dunedin School of Medicine, University of Otago, Dunedin, New Zealand

^h Department of Surgery, John A. Burns School of Medicine, University of Hawaii, Honolulu, USA

ⁱ British Vein Institute, London, UK

Submitted 5 November 2010; accepted 17 March 2011

Available online 6 May 2011

KEYWORDS

Consensus document;
Duplex ultrasonography;
Varicose veins;
Surgical treatment;
Endovenous treatment;
Follow-up

Abstract Objectives: Duplex ultrasound has become the reference standard in assessing the morphology and haemodynamics of the lower limb veins. The project described in this article was an initiative of the Union Internationale de Phlébologie (UIP). The aim was to obtain a consensus of international experts on the methodology and terminology to be used for assessment after treatment of incompetent superficial and perforating veins in the lower limb by ultrasound imaging.

Design: The study design was consensus meetings leading to a consensus document.

Methods: The UIP invited group submitted relevant literature references and written contributions concerning the methodology, terminology and value of duplex imaging after treatment. The authors prepared a draft document that was circulated to a larger group of experts and revised according to the comments received. Eventually, all participants agreed upon the final version of the article.

CME To access continuing medical education questions on this paper, please go to www.vasculeducation.com and click on 'CME'

* Corresponding author. M. De Maeseneer, Department of Dermatology, Erasmus Medical Centre, PB 2040, 3000 CA Rotterdam, The Netherlands. Tel.: +31 10 7033798; fax: +31 10 7033822.

E-mail addresses: marianne.demaeseneer@ua.ac.be, m.demaeseneer@erasmusmc.nl (M. De Maeseneer).

Results: Formal analysis of the results of interventions for varicose veins relies on adequate preoperative assessment and a careful description of the procedure employed. The timing of investigations of outcome should be classified as immediate (1–4 weeks), short-term (1 year), midterm (2–3 years) and long-term (5 years or more). The examination should employ standard methodology and formally described variables, which can be tailored to the intervention that was undertaken. The experts have made detailed recommendations concerning the methods to be used for duplex ultrasound examination and reporting after various treatments for varicose veins, including novel treatments under scientific study.

Conclusions: Duplex ultrasonography is a fundamental component of the investigation of the lower limb venous system after treatment for varicose veins.

© 2011 European Society for Vascular Surgery. Published by Elsevier Ltd. All rights reserved.

Duplex ultrasound (DUS) is the most frequently used investigation to evaluate the outcome of treatments for chronic venous disease (CVD). Endovenous ablation (EVA) of a vein using laser or radiofrequency energy, or ultrasound-guided injection of sclerosant foam may all be evaluated using this technique. DUS imaging is the ideal non-invasive method for follow-up, as it provides anatomical and haemodynamic information about the treated veins.^{1–9} DUS can detect the early stages of recurrent varicose veins, before they become apparent clinically.¹⁰ Serial DUS imaging can not only help to understand the clinical evolution of the individual patient after treatment for CVD, but also has the potential to increase the general knowledge of events leading to clinical recurrence. Thus, long-term follow-up using DUS extends the understanding of the natural evolution of varicose vein disease.

Widely different DUS criteria have been used to assess the outcome of treatment for venous disease; often, there is very little information about the preoperative morphological and haemodynamic condition of veins. Currently, there is no systematic agreement from phlebology or vascular societies on how DUS imaging is best performed, or interpreted for follow-up. Standardisation of follow-up imaging and reporting would reduce the confusion and give better clarity to the end points of treatment.^{11–13} The aim of this document is to summarise best practice for venous DUS examination of the legs after treatment, derived partly from the (limited) published evidence, and also agreed upon by an expert group that regularly uses this technology.

Methodology

The Union Internationale de Phlébologie (UIP) invited a group of international experts in the field of DUS investigation with MDM as the Chair. They were invited to submit relevant literature references and written contributions concerning the methodology and value of DUS imaging after treatment. Personal expert opinions were sought, which did not necessarily reflect policies of scientific or medical societies to which the individuals were affiliated. This process was not intended to form a systematic review of the literature, but to provide evidential support to the consensus recommendations made in the final document.

Consensus meetings began with interested experts at the Venous Forum of the Royal Society of Medicine in Manchester, November 2007, and subsequent meetings

were held with smaller subgroups looking at specific areas. The aim was to define a systematic method of DUS imaging and reporting, which was simple to use and could be employed to audit the outcome of various treatments for varicose veins, including novel treatments under scientific study. The authors acknowledge that many of the statements only reflect the opinion of the consensus group rather than being supported by published scientific evidence. This reflects the fact that existing publications have used widely varying criteria for both methods and interpretation of DUS imaging after treatment.

Only the specifics of DUS imaging for the evaluation of treatment are considered. However, it is critical that these go together with the essentials of good clinical and patient-focussed evaluation. The tools for these have been published previously, such as Clinical, etiologic, anatomic and pathophysiologic data (CEAP), Venous Clinical Severity Score (VCSS) and other generic and condition-specific (e.g., the Aberdeen Varicose Vein Symptom Severity score, AVVSS) quality of life (QoL) assessments.^{14–19} The relationship of these tools to specific DUS appearances is unfortunately not yet established. Further work is required, which should be stimulated by this consensus document. It is anticipated that, in future studies of global outcome following different treatment strategies, the present document should not be used in isolation, but should be used in conjunction with similar existing or newly produced consensus documents on clinical reporting.

While this consensus deals primarily with the DUS evaluation after treatment, it cannot ignore the state of the pre-treatment evaluation. Anatomical and haemodynamic characteristics of superficial venous insufficiency often vary widely between patients. Therefore, clear recording of pre-treatment clinical features and DUS findings is essential, as they may influence the interpretation of post-treatment results and expectations. The detailed documentation of the type of treatment employed is also important for the understanding of subsequent DUS findings. For instance, the length of the incompetent segment of the great saphenous vein (GSV) that was treated should be recorded, and the anterior accessory saphenous vein (AASV) should be distinguished clearly from the GSV. These pre-treatment details are unfortunately not always available, which may significantly reduce the value of DUS-based follow-up. The previous consensus document on recurrent veins after surgery (at that time called 'REVAS'), which integrated clinical and DUS findings, reported that, for the majority of patients, little information was

available regarding the type of procedure undertaken, or the date it was done.^{20,21} In the recently published Vein-TERM consensus document, the acronym PREVAIT (PREsence of Varices (residual or recurrent) After InTervention) has been introduced, referring to clinical scenarios where varices cannot definitely be classified as recurrent or residual.²² In case of PREVAIT, interpretation of DUS findings is always difficult and potentially ambiguous. The present consensus document seeks to address this by insisting on the recording of all useful prior information, and on the requirement for prospective DUS evaluation for adequate interpretation.

DUS Imaging before Treatment

The basic principles of venous DUS investigation for varicose veins have been described in a previous UIP Consensus Document, and these remain applicable.⁸ The ultrasound anatomy of the superficial and perforating veins has also been described in a UIP Consensus Document.⁹ Since these publications, new treatments have emerged for which additional descriptive features are required, and these are included in the present document.

The minimum requirements for pre-treatment DUS assessment are described in Table 1. Other features may be recorded, but should not be at the expense of the minimum data set.

To assess the involvement of any pathology of the deep venous system, scanning the deep veins and looking for patency and the presence/absence of reflux is an important part of the evaluation of a patient before treatment of varicose veins. Some patients present with superficial vein disease due to underlying primary deep venous incompetence or secondary (post-thrombotic) obstruction and/or reflux. This will obviously influence the long-term outcome of treatment of superficial or perforating vein incompetence adversely and should therefore be included in the initial assessment as well as in further follow-up. In

patients with a previous history of deep vein thrombosis, the deep venous system should be examined in both the lying and standing position to check for residual obstruction and reflux, respectively, as described previously.⁸

When assessing superficial veins, patients should be examined in the standing position where possible, to standardise measurements of venous diameter and reflux. If obesity or other cardio-respiratory conditions make this impractical, it is vital that any follow-up imaging is done in the same position.

Diameter measurement

This should be performed in transverse view and the outer diameter should be measured (including the vein wall) to compare this with the postoperative diameter after endovenous ablation. For the GSV trunk, diameter measurements of the incompetent segment should be made 3 cm below the saphenofemoral junction (SFJ), at mid-thigh level, at the knee and also at mid-calf level. The saphenous trunk should be measured at a site where there is no focal (or aneurysmal) dilation of the trunk.²³ Similarly, the AASV should be measured 3 cm below the SFJ and at mid-thigh (if the trunk exists at this level). For the small saphenous vein (SSV), the trunk should be measured 3 cm below the saphenopopliteal junction (SPJ) where the pre-terminal valve, if present, is located. A mid-calf measurement should also be made.

Assessment of reflux

The presence of flow in any part of the arterial or venous system assumes the existence of a pressure gradient; nor can venous reflux exist without a pressure gradient. In an individual standing still and breathing normally, flow in both competent and incompetent veins is very slow but remains antegrade. To detect the presence of reflux, it is necessary to create a pressure gradient in the venous system. This is usually achieved either by a Valsalva manoeuvre, which creates high pressure in the venous system, or by compression–release (calf squeeze), which creates a low pressure in the venous system during release.

There is a lack of standardisation in reflux assessment; the duration of reflux in seconds (reflux time) is used most commonly. Reflux is usually defined as retrograde flow lasting for more than 0.5 s, whereas less than 0.5 s is defined as normal or no reflux.⁸ However, the reflux time can only be used to distinguish between a competent and incompetent vein segment and is therefore a qualitative evaluation. Peak reflux velocity (m s^{-1}) and the rate of reflux (ml min^{-1}) may be used to provide a quantitative evaluation of reflux.²⁴ Manual limb compression by calf squeeze cannot be standardised; hence, is not ideal for quantitative measurement of reflux. To obtain more reproducible results, two standardised methods for testing reflux have been evaluated in healthy subjects and patients with venous disease:

1. Standardised Valsalva manoeuvre: This consists of forced exhalation into a special tube system that measures expiratory pressure (Jeanneret–Valsalva device, Eisenhut® – VET Ag, Allschwil, Switzerland).

Table 1 Preoperative duplex imaging.

1. <i>Deep veins</i> : assessment for patency and reflux
- common femoral vein (CFV)
- popliteal vein
2. <i>Junctions</i> : assessment for reflux (terminal valve/pre-terminal valve)
- saphenofemoral junction (SFJ)
- saphenopopliteal junction (SPJ)
3. <i>Main trunks</i> : diameter measurement and assessment of reflux (in the saphenous compartment):
- great saphenous vein (GSV)
- anterior accessory saphenous vein (AASV)
- posterior accessory saphenous vein (PASV)
- small saphenous vein (SSV)
- thigh extension of SSV/Giacomini vein
4. <i>Tributaries</i> : if incompetent
5. <i>Non-saphenous veins</i> : if incompetent
6. <i>Perforating veins</i> : diameter measurement and assessment of reflux

A pressure of 30 mm Hg should be established within 0.5 s and held for at least 3 s, to avoid false positive prolonged reflux due to a lack of transvalvular pressure gradient. This standardised Valsalva manoeuvre is highly reproducible in the proximal leg veins with coefficients of variation ranging from 10.4% to 13% for reflux time and from 11.5% to 16.4% for peak reflux velocity.²⁵ The Valsalva manoeuvre, however, cannot be used to assess reflux in veins distal to competent valves.

2. Standardised cuff inflation–deflation method: An automatic or manual pedal cuff inflator produces rapid inflation and deflation of cuffs placed at different levels on the leg.²⁶

In most countries, these standardised methods are not used routinely. Most assessors use a Valsalva manoeuvre, together with calf compression–release to characterise venous incompetence. At the level of the SFJ, the combination of these two manoeuvres is essential to assess the state of the terminal valve (TV) of the GSV. The TV and the pre-terminal valve (PTV) of the GSV should both be identified, to distinguish between incompetence of both valves or only one of the two.⁹ The assessment of TV and PTV haemodynamics is best achieved through the combined use of colour flow and Doppler modalities, with the sample volume placed above each valve to test for the presence or absence of reflux (at common femoral vein level to test the TV and at the most proximal part of the GSV to test the PTV).

In the groin, the lymph node area should be studied, particularly in patients with recurrent varicose veins. Reflux in a lymph node vein network (LNVN) should be sought, tracing it upwards looking for a connection with the common femoral vein or pelvic veins, and downwards looking for a connection with GSV, AASV or other varicose veins.²⁷

In multiparous women, a leash of tortuous veins may also be seen within the saphenous compartment or superficial to the SFJ in the thigh. These veins mostly commonly connect with the abdominal–pelvic venous network and reflux may be elicited during Valsalva or calf release. They may or may not connect with the main trunk of the GSV or AASV. DUS imaging may reveal reflux in one or more of these visible veins.²⁸

Details of Treatment or Procedure

Sufficient details of any treatment should be documented to permit informed DUS follow-up. These are described as minimum requirements in Table 2. Additional details can be included, as required, to fulfil the requirements of specific studies.

Timing of Post-treatment DUS Evaluation

There are several purposes for DUS follow-up. These include assessment of adequacy of the initial treatment and of complications, identifying possible need for further intervention, identifying the evolution of deterioration and recurrence, carrying out research into treatment mechanisms and outcomes and, in due course, to identify surrogate end points of long-term outcomes. The intended

Table 2 Details of treatment.

1. Treatment of junction (SFJ, SPJ)
- none
- flush SFJ ligation
- lower ligation with preservation of the junction itself
- repeat ligation of SFJ or alternative procedure for recurrence
- adjunctive procedures: closing opening in cribriform fascia, prosthetic patch on SFJ/SPJ
2. Treatment of main trunk (GSV, AASV, SSV)
- none
- endovenous laser ablation
- radiofrequency ablation
- (ultrasound-guided) foam/liquid sclerotherapy
- stripping with vein stripper (invaginated or not) or cryostripping
- length/site of treated trunk(s)
3. Treatment of tributaries
- none
- phlebectomies
- foam/liquid sclerotherapy
- endovenous laser treatment
- delayed phlebectomies or sclerotherapy
4. Treatment of perforating veins
- none
- phlebectomy and epifascial ligation
- foam/liquid sclerotherapy
- radiofrequency ablation
- endovenous laser ablation
- subfascial ligation (SEPS)
- site of treated perforating vein(s)

purpose, as well as the cost and the feasibility of inviting patients to re-attend, will influence the frequency of repeated assessments.

Immediate: 1–4 weeks after treatment

This is indicated:

- (1) For single 'one-stop' treatments, such as surgery or endovenous thermal ablation, where it is desirable to know whether the intervention has achieved the intended immediate goal. Without this, it is impossible to determine whether recurrence, should it occur, is due to inadequate therapy (for instance, residual incompetent GSV trunk after stripping). The presence of post-treatment deep vein thrombosis should also be assessed.^{29–33}
- (2) As part of sequential treatments, such as staged (foam) sclerotherapy, combined procedures (phlebectomy + sclerotherapy), CHIVA (Cure Conservatrice et Hémodynamique de l'Insuffisance Veineuse en Ambulatoire) technique,³⁴ etc.

Late follow-up

For most patients, further DUS evaluation will be required for suspected recurrence. Repeat assessment is essential for

longer term prospective evaluation intended for the purposes of research or clinical audit. For these purposes, recommended follow-up is:

Short-term. Assessment should be performed 1 year after the intervention to elicit ultrasound features that may predict longer term outcome after treatment of GSV, AASV or SSV. A vein obliterated at 1 year after endovenous thermal ablation will usually remain so for at least 3–5 years in most cases.^{35,36} After surgery, 1-year assessment detects newly developed incompetent veins (due to neovascularisation or dilation of pre-existing veins) in the treated areas.^{37–41} The finding of new incompetent veins in the groin, which are larger than 4 mm in diameter, 1 year after SFJ ligation results in a 100% probability of developing a clinical groin recurrence and visible recurrent veins in the thigh after 5 years.¹⁰

Mid-term. This should be considered after 2–3 years to confirm the earlier changes; but, this interval after treatment is still too short for conclusions on long-term outcomes.^{6,41–43}

Long-term. This should be performed at least 5 years after treatment.^{6,36,44–47} This interval is long enough for the development of clinical recurrence, which may have arisen as a consequence of the incompetent veins detected by the short- or midterm DUS scan.^{6,10,36}

In addition to DUS imaging, clinical findings should be evaluated using VCSS (or an alternative clinical score) and QoL assessment at the same time intervals.¹²

If possible, a late DUS assessment can be performed after 10 years or more.^{46,48–50} The longest reported DUS follow-up after varicose vein surgery was done 34 years after treatment, but, for obvious reasons, this was not a prospective study.⁵¹ In practice, it is not easy to achieve long-term follow-up in patients treated for varicose veins.

DUS Imaging after Surgical or Endovenous Treatment of Varicose Veins

Standardised preoperative DUS assessment and clinical evaluation (see above) should combine a written report with a diagrammatic representation of the data as an adequate basis for further DUS comparisons.⁸

A. DUS imaging after surgical treatment

Following surgical treatment, DUS imaging should focus on the saphenous junctions (SFJ and SPJ) and keep track of the stripped saphenous trunk (GSV, AASV or SSV). Sites of previous perforator ligation should also be investigated. Recurrent varices should be assessed as either:

- (a) veins with reflux following a Valsalva or compression–release manoeuvre, where a change of compartment or escape point (connection with the deep veins) exists; and
- (b) veins where the reflux is not linked to an escape point but is generated by the filling of the incompetent vein by its tributaries. In these veins, a Valsalva manoeuvre will not elicit reflux, whereas a compression–release manoeuvre will show reflux in the relaxation phase.

All ‘escape points’ should be documented, where possible; in many cases of recurrent veins, no clear communication with deep veins can be identified by DUS examination.^{20,21,45}

DUS of the SFJ

The SFJ is a common site of recurrence following surgery.^{21,52} After classic SFJ disconnection (flush saphenofemoral ligation and ligature-section of the related SFJ tributaries), also called crossotomy or high ligation, a number of features may be identified:

- **Normal:** This shows the common femoral vein (CFV) without any residual segment of GSV or any incompetent superficial vein in the groin, and the GSV terminal valve is no longer visible.
- **Neovascularisation or groin varicose network:** Neovascularisation is defined as new blood vessel formation (angiogenesis) occurring in abnormal tissue or in an abnormal position, which is a histological diagnosis. However, in the Vein-term consensus document, neovascularisation has been accepted as a venous term, defined as: “presence of multiple new, small tortuous veins in anatomic proximity to a previous venous intervention.”²² The consensus group has retained the term ‘neovascularisation’ to describe the presence of new veins situated at the site of the previous SFJ or SPJ ligation (Fig. 1(a) and (b)).^{7,53–56} These veins may be newly formed or can arise from dilation of existing groin veins that were invisible on DUS before the operation. They can be observed in the short-term, mid-term or only at long-term follow-up.⁴⁶ The terms ‘neoreflux’, ‘angiodysplasia’ and ‘cavernoma’ should not be used anymore to avoid confusion with other conditions.^{56–58} An alternative term for neovascularisation at the SFJ could be ‘groin varicose network’, which is a neutral descriptive term for the newly visible veins, without suggesting any underlying pathophysiology. If tortuous veins are seen in connection with a lymph node, they are properly described as LNVN and these findings may or may not be part of the neovascularisation pattern.⁵⁸ Neovascular veins are usually tortuous. They may exhibit reflux with a Valsalva manoeuvre, and/or during release after calf compression. Some neovascular veins have a normally directed flow without reflux. With time, these veins may establish a clear connection with superficial tributaries or with an LNVN, or with a retained saphenous trunk (which was not stripped intentionally or unintentionally) or with other veins present in the saphenous compartment after stripping, and become larger. The largest diameter of the vein(s) of the vascular network in the groin should be measured and the veins should be checked for presence or absence of reflux during calf compression–release and/or Valsalva manoeuvre. If these veins exhibit reflux with a Valsalva manoeuvre, the escape point usually lies in the SFJ area. In some cases, there is a connection with incompetent pelvic veins. If venous reflux is detected in neovascular veins only during the calf release phase (and not during Valsalva manoeuvre), this suggests that they fill from the subcutaneous abdominal venous




Figure 1 (a) Network of tiny refluxing veins originating from the common femoral vein (CFV) in the high ligation area suggesting angiogenesis (neovascularisation). (b) Network of large refluxing veins originating from the common femoral vein in the high ligation area without residual stump neither terminal valve.

network towards the groin, thigh and leg veins, without a direct communication (escape point) with the deep veins.

- *Residual stump*: If GSV ligation has been performed at a distance from the CFV (low ligation) instead of flush with the SFJ, a residual stump may be seen. The terminal valve can usually be seen with one or more residual SFJ tributaries. The most common pattern of recurrent veins is venous reflux via the SFJ, although this is commonly associated with reflux in residual tributaries. Reflux from the residual stump can connect with a residual AASV or other varices (Fig. 2). Where the terminal valve is competent, the stump receives inflow from its tributaries that drain normally into the SFJ. This pattern of flow is usually seen after successful EVA, but may also be found after selective stripping of the GSV trunk without a flush ligation, preserving the saphenofemoral confluence.^{59,60} In prospective studies, the diameter of a residual stump should be measured and reflux in the stump or any of its connections should be assessed on DUS.




Figure 2 Network of large refluxing veins originating from the common femoral vein (CFV) through a residual GSV stump with the incompetent terminal valve still visible (arrow).

DUS of the above-knee GSV and of the AASV

After stripping of the above-knee GSV, the saphenous compartment should be examined, to see whether the GSV really is absent. If the GSV trunk is still completely or partially present within its 'saphenous eye', the diameter and length of the residual GSV segment should be recorded, and reflux assessed using a calf compression–release manoeuvre.

In the case of CHIVA, the persistence of the GSV trunk with retrograde flow is part of the treatment itself, which aims to redirect the saphenous blood flow towards competent re-entry perforators in the thigh or calf.⁵⁰

In patients who have had surgical stripping, revascularisation of the strip-track has been observed, with the presence of multiple convoluted channels in the track of the previously stripped GSV.⁶¹ It would be better to describe this only as 'multiple venous channels in the saphenous compartment' (see the primary varicose network described earlier).²⁸ The DUS image is typical in transverse as well as in longitudinal view (Fig. 3). Reflux can




Figure 3 Early postoperative scan after GSV stripping showing haematoma still visible in the strip track (arrows) containing a network of multiple tiny refluxing veins.

be found in these venous channels and they may connect with clinically obvious recurrent varicose veins. Reflux may persist in the varicose network after stripping the GSV, or the varicose network may dilate and become incompetent postoperatively.

As the AASV is often involved in recurrence after GSV stripping, this vein should be examined along its course (usually from mid-thigh to the groin), including diameter measurement. In particular, connections with any possible escape point(s) usually located more cranially should be examined, for example, a residual stump, refluxing pelvic veins, LNVN, perforators, etc.

Much less common is recurrence in the posterior accessory saphenous vein (PASV) (sometimes in connection with a Giacomini vein). Typically in multiparous women, incompetent abdominal–pelvic veins can connect directly with residual GSV segments, or superficial tributaries in the thigh after surgery, although these may have been present before surgery.²⁸

DUS of the below-knee GSV

The below-knee GSV is studied in its saphenous compartment. Residual reflux may be present in part or all of the saphenous trunk below the knee, and the diameter should be measured. GSV reflux within the saphenous compartment below the knee may re-enter the deep venous system through a distal re-entry perforator, without a connection to any superficial varicose veins.

DUS of the SPJ and SSV

After SSV surgery at the level of the SPJ, DUS findings are comparable to those at the SFJ after GSV surgery. However, it is important to have details about the preoperative anatomical and haemodynamic situation, as well as the procedure performed, as this will assist interpretation of the findings at the SPJ. Variations in SPJ anatomy are common and haemodynamics of the popliteal fossa are complex. The varying level of the SPJ in relation to the popliteal skin crease, the potential presence of a thigh extension of SSV or a Giacomini vein, the possible junction of the SSV with one of the gastrocnemius veins before joining the popliteal vein, the presence of a popliteal fossa perforating vein and all the different haemodynamic situations related to these anatomical variations has been described in detail in the two previous UIP Consensus Documents on DUS investigation (including an illustrative diagram).^{8,9}

DUS imaging may highlight:

- *normal postoperative findings*. It is essential to specify, if ligation has been performed flush at the level of the popliteal vein, or at the confluence with one or more gastrocnemius veins; in the latter case, a residual stump has been left intentionally, since it represents a common track between gastrocnemius vein(s) and the proximal SSV.
- *neovascularisation, or popliteal fossa varicose network* with, or without reflux. Comparable to the situation in the groin, the neovascular veins may connect directly to the popliteal vein, or they may connect with incompetent veins in the posterior thigh (e.g., Giacomini vein, thigh extension of the SSV, pelvic or gluteal

veins, sciatic nerve varices and sciatic veins). Typically, reflux is most obvious during calf release (diastolic phase). However, in a few cases, recurrent reflux within the varicose network in the popliteal fossa and/or in the residual stump may be elicited during calf compression or contraction (systolic phase), with, or without reflux during calf release. This specific haemodynamic pattern is exceptional and may indicate impaired outflow in the popliteal and/or femoral vein due to anatomical or functional abnormalities.⁸

- *residual stump*, with or without reflux. An incompetent residual stump is frequently seen after SSV surgery. This may arise from the great variation in level of the SPJ.⁹ Unfortunately, data concerning the outcome of SSV surgery including mid- or long-term follow-up are extremely scarce and very few studies have focussed on the clinical recurrence rate and typical DUS appearances.^{37,49,62–65}

Recurrent varicose veins after SSV surgery may be related to other kinds of anatomical and haemodynamic patterns:^{9,66} (pre- or) postoperative incompetence of the popliteal fossa perforating vein, which is a tortuous vein, usually located laterally from the SPJ (Fig. 4(a) and (b)),




Figure 4 (a) A popliteal fossa perforating vein can be recognised from its typical location in front of the lateral condyle of the femur (arrows). (b) Colour duplex image shows reflux in this tortuous vein.

(pre- or) postoperative gastrocnemius vein incompetence or (pre- or) postoperative popliteal vein incompetence. Incompetent proximal veins (e.g., pelvic or gluteal veins, sciatic nerve varices (Fig. 5) and sciatic veins) can connect directly with residual SSV segments, and are frequently seen in women with pelvic varicocoele.

DUS imaging of perforating veins

Perforating veins are described according to their location above or below the knee, medial, anterior, lateral or posterior as described in the UIP consensus document.⁹ Perforating vein incompetence is defined as a perforator that allows outward flow of >0.5 s during calf relaxation or release after distal compression.^{8,22} The fate of perforating veins after surgical treatment has been studied extensively in a large prospective study, which included DUS assessment at 6 months, 1 year and 3 years.⁶⁷ This study showed that recurrent perforator incompetence after surgery is far more common than previously recognised; correct adjustment of DUS equipment is required to highlight low-velocity reflux during calf relaxation or release, which may occur in small perforators. In primary venous insufficiency, perforators that transmit part or the totality of the reflux from the superficial into the deep venous system are described as re-entry perforators.^{8,68} After saphenous stripping and phlebectomy, they mainly show normal inward flow and a diameter reduction at short- and mid-term follow-up, though long-term data are still lacking.^{69,70}

DUS imaging of tributaries and of non-saphenous veins

During DUS follow-up, the final report should include an assessment of the main incompetent tributaries, the

intersaphenous veins and the non-saphenous veins (e.g., the lateral venous system).⁹ On the back of the thigh, varicose veins may receive reflux from a gluteal plexus or vulvar/perineal veins, and are often related to pelvic vein incompetence. By carefully tracing these veins upwards while eliciting reflux with a distal compression–release manoeuvre, it is possible to detect the related pelvic vein reflux. If clinically indicated, in selected patients, this may be further evaluated by additional DUS or venographic investigation of pelvic veins.

B. DUS Imaging after Physical or Chemical EVA (Laser, Radiofrequency or Foam Sclerotherapy)

EVA using laser energy, radiofrequency-generated thermal energy or a chemical sclerosant (most frequently as foam) has become increasingly popular for the treatment of varicose veins. Wide varying criteria have been used to classify the outcome after EVA.⁷¹ A joint statement ‘Recommended reporting standards for EVA for the treatment of venous insufficiency’ has been published recently.¹² DUS efficacy criteria after foam sclerotherapy were discussed during a European Consensus Meeting in Tegernsee, Germany in 2006, and it was suggested these criteria could be suitable for other EVAs too.¹³

To evaluate EVA of the GSV, SSV or AASV, DUS investigation should focus on both the saphenous junctions (SFJ and SPJ) and on the treated trunk. If additional surgery of tributaries and/or of perforating veins has been performed, DUS findings will be reported as mentioned above. DUS imaging will describe the morphological and haemodynamic situation at the same follow-up intervals as those for evaluation of clinical outcome.

Thermal methods for EVA aim to obliterate the whole vein length between the site of introduction of the fibre or catheter and the incompetent junction. During radiofrequency ablation (RFA), the catheter tip is usually positioned 1 cm distally from the ostium of the superficial epigastric vein (2 cm from the SFJ).^{72–74} For endovenous laser ablation (EVLA), the tip of the laser fibre is advanced to within 0.5 and 3 cm from the SFJ.^{75–77} Similarly, for SSV treatment, the catheter or fibre tip is positioned 1–2 cm below the SPJ.^{78–80} In ultrasound-guided foam sclerotherapy (UGFS) or catheter-directed foam sclerotherapy, an adequate volume of foam is delivered to obliterate the target vein, usually in one or two sessions.^{13,43,81–83}

DUS of the SFJ

After endovenous treatment of the GSV with laser, RFA or foam, the terminal part of the GSV usually remains open and permits drainage of one or more tributaries of the SFJ.

- *Morphology: obliteration or patency?* The presence of a patent terminal portion of the GSV is considered to be a normal finding after endovenous treatment, if it is <3 cm in length. When only the distal incompetent part of the GSV has been treated, the competent proximal section will exceed 3 cm. The proximal section of the GSV may receive inflow from a Giacomini vein, pudendal veins and other tributaries.⁷² At the immediate DUS scan (1–4 weeks after treatment), thrombus




Figure 5 Sciatic nerve varices (posterior thigh view).




Figure 6 (a) Thrombus extension into the CFV (arrows) after thermal EVA (thrombus has developed in the GSV stump). (b) Thrombus extension into the CFV after foam sclerotherapy—thrombus has developed along the anterior wall of the GSV and CFV (arrows).

extension from the GSV into the CFV should be sought (Fig. 6(a) and (b)). Protrusion of a thrombus into the lumen of the CFV is very unusual (<1%) and should always be considered a pathological finding.^{11,30–33,84,85} The extent of the thrombus should be assessed to assist in selecting appropriate treatment and monitoring.⁸⁶

- *Haemodynamics: presence or absence of reflux.* Detection of low-velocity reflux requires proper adjustment of Doppler and colour modules,⁸ and is of great importance in assessment after EVA, where vein compliance is usually reduced (because of diameter reduction or disappearance); tiny vessels may show reflux. To elicit venous reflux, a sufficiently high compliance is necessary, while the energy/pressure gradient is the second component.⁸⁷ After EVA, DUS usually shows a patent SFJ or SPJ with obliteration of the upper GSV/AASV or SSV. This obliteration dramatically reduces the compliance of the remaining few centimetres of patent vein. As a result, there is abolition of the energy gradient needed to generate reflux from the CFV through the SFJ, although valvular

incompetence may still exist. Reflux will then not be detectable during Valsalva and compression–release manoeuvres in the proximal part of the GSV. The long-term haemodynamic fate of the residual saphenous stump after EVA is debated and its course should be monitored during DUS follow-up. Persistence or reappearance of reflux at the SFJ and/or at the level of the saphenous stump is always considered pathological. Any varicose veins in connection with this vessel should be imaged; infrequently, this consists of an incompetent non-obliterated saphenous trunk, but more often recurrent thigh varicosities connect with the incompetent part of the GSV in the groin, either through an incompetent AASV or PASV, or directly. In some cases, reflux can be provoked by a Valsalva manoeuvre or after thigh compression (more easily than after calf compression) in one or more tiny veins in the groin lymph node area and this finding mimics neo-vascularisation after saphenous stripping. Usually, no connection with any visible varicose veins is detectable at short-term follow-up. After EVA, the incidence of this DUS pattern is very low (0–1%).^{88,89} The possible future clinical significance of this finding is not clear yet and the longest reported follow-up is limited to 5 years.³⁶

DUS of the above-knee GSV and of the AASV

Obliteration of the trunk can usually be observed on completion of the procedure, but it usually takes 6–12 months before complete ultrasonographic disappearance is achieved.^{72,89}

- *Morphology: obliteration or patency?* Treated veins (GSV and/or AASV) should be assessed by their compressibility, as well as using colour flow. It is standard practice to treat the whole above-knee GSV with EVLA and RFA and this vein should be obliterated in its entire course. Foam sclerotherapy may be less targeted, but, in most cases, a similar procedure and outcome is expected. However, the sonographic appearance of the obliterated vein will depend on the stage of post-treatment evolution. In the initial stages, the vein behaves differently depending on whether it has been treated with EVLA, RFA or foam.^{90,91} Subsequent obliteration is characterised by progressive reduction in vein diameter, often in an inhomogeneous way over the length of the treated vein. In the final stage of fibrotic transformation, the vein may disappear completely on DUS imaging, or is transformed into a fibrous cord that may be visible as a hyperechogenic tract in the saphenous compartment. The time course of this whole evolution varies between patients.⁸⁹ The outer diameter of the obliterated vein can be measured, preferably on a transverse section, and the residual inner lumen of the visible vein can be assessed in case of partial or complete patency. Sometimes, an obliterated vein can be seen with an inhomogeneous content characterised by varying echogenicity, without flow or reflux, but with a lumen that can be partially compressed. The latter may be due to the presence of blood or thrombus in the lumen in an early stage.

- **Haemodynamics: presence or absence of reflux.** The presence of reflux following a Valsalva or compression–release manoeuvre is pathological. This reflux may be present over the whole length of the treated vein or may be segmental and the possible escape points should be described (SFJ, perforating vein, refluxing pelvic veins, etc.). Reflux distribution in the saphenous trunk and/or tributaries, together with the re-entry or refluxing perforators should be noted. Whether the reflux is due to primary failure of EVA treatment or re-canalisation after initial successful obliteration (usually within 6 months) can only be determined by performing serial DUS imaging. Antegrade flow without reflux can sometimes be demonstrated in a partially or completely patent residual GSV trunk. This may result from reduced vein size and the obliteration of re-entry points. No conclusive data are available to help interpret the significance of this finding, which might represent a good pathophysiological result, as the reflux is abolished.¹³ The most important morphological and haemodynamic DUS findings after EVA of a saphenous trunk are summarised in Table 3.
- **Additional DUS evaluation above the knee.** As the AASV may be involved in recurrence after GSV ablation, this vein should always be examined along its course, and the presence or absence of reflux should be assessed.⁷⁶ Vice versa, after previous AASV ablation, the GSV main trunk should be examined at follow-up DUS. Although the PASV (and Giacomini vein) can also play a role in recurrence, this is far less frequent.

DUS of the below-knee GSV

Although it is still a controversial issue,⁹² usually the GSV is ablated just to the level of the knee, even if the below-knee segment is also incompetent. Immediately after the

Table 3 Main morphological and haemodynamic duplex ultrasound findings after endovenous ablation of a saphenous trunk.

Morphology:

- (a) obliteration of the vein: total incompressibility and absence of colour flow; the vein can be classified as visible or nonvisible
- (b) partial patency of the vein: partial compressibility and presence of colour flow in a part of the lumen
- (c) total patency of the vein: complete compressibility and presence of colour flow in the totality of the lumen
- (d) outer diameter of the residual vein
- (e) inner diameter of the residual lumen in case of partial or total patency
- (f) segmental obliteration/patency: length of the obliterated, partially or completely patent segment(s)

Haemodynamics:

- (a) absence of flow with both principal manoeuvres (Valsalva, compression/release)
- (b) antegrade flow during compression manoeuvre
- (c) retrograde flow >0.5 s during one or both manoeuvres

EVA procedure, the GSV below the knee may remain competent or incompetent, or it may become incompetent again later at DUS after an interval. Partial or complete thrombosis of the below-knee GSV may occur after ablation of the above-knee GSV.

DUS of the SPJ and SSV

Sonographic changes after endovenous treatment of the SSV have been described less frequently than after GSV treatment. Changes in the SPJ and SSV seem to be similar to those of the SFJ and GSV, and should therefore be reported in the same way. At the SPJ, proximal tributaries may remain patent, particularly any extension of the SSV into the thigh or the Giacomini vein. Taking into account the variable level of termination of the SSV, the length of patent SSV considered acceptable after treatment should take the level of the SPJ as reference, rather than the skin crease of the popliteal fossa. After EVA of the SSV, obliteration of the treated SSV segment should be assessed and the residual distal SSV segment should be tested for reflux.

DUS imaging of perforating veins

Incompetent perforating veins can be treated with laser or radiofrequency using a technique-limiting thermal ablation to a short vein segment, or with foam sclerotherapy. The efficacy of all these treatments can be evaluated with DUS imaging, illustrating the obliteration of the treated perforating vein and abolition of outward flow during the release (diastolic) phase of the compression–release manoeuvre.^{8,93,94} Perforating veins that remain patent should be reassessed to compare the normal inward flow and pathological outward flow, with the aim of determining the net flow. The latter is obtained by comparing mutually pulsed Doppler curves of inward and outward flow during distal compression and release.^{8,68} The course of the reflux should be traced whenever it extends from the perforating vein into any saphenous or tributary vein.

Standardisation of results after EVA in view of scientific studies

The authors propose a classification (Table 4) allowing precise analysis of the results after EVA, to enable comparison of different techniques and to introduce

Table 4 Proposed classification of duplex findings at the junction (J) and at the treated trunk (T) after endovenous ablation.

J: for SFJ or SPJ:

- J0: no patent stump
- J1, J2, J3, J4 etc.: junction with patent stump of 1, 2, 3, 4 cm etc.

R+: reflux; R-: no reflux

T: for GSV/AASV/SSV trunk:

- Ti: invisible trunk
- To: obliterated trunk (diameter: mm)
- Tp: completely or partially patent trunk (diameter: mm)
- To/Tp or Tp/To: segmental obliteration/patency or patency/obliteration (length of patent segment: ... cm; diameter of residual lumen: ... mm)

R+: reflux; R-: no reflux

a standard DUS assessment of outcome following any EVA procedure. The situation at the level of the junction (J) and the treated trunk (T) is recorded. Reflux is reported as present (R+) or absent (R-). DUS findings at the level of the SFJ or SPJ are described as J0, J1, J2, J3, J4, etc., according to the absence of any visible stump (J0) or the presence of a stump with a patent length of, respectively, 1, 2, 3, 4 cm etc. If reflux is present, the site of connecting refluxing tributaries should be specified. DUS findings at the level of the treated saphenous trunk are reported as Ti (invisible), To (obliterated) or Tp (patent) or a combination of the latter in case of segmental obliteration. The presence or absence of reflux is reported as R+ or R-. If reflux is detected, the site of refluxing tributaries should be described. Measurement of diameter and length of the obliterated and/or patent segment of the saphenous trunk are described using the same protocol. When the lumen of the treated vein is partially obliterated in a segment of the treated vein or in its total treated length, the diameter of the residual lumen can be measured as well (Table 4).

The Role of DUS Investigation in Clinical Studies

Clinical outcome studies are designed to measure the impact of therapy on clinically relevant variables, such as patient satisfaction, quality of life and relief of symptoms. They also examine clinical signs, such as the presence of varicose veins, oedema and skin changes including venous ulceration. DUS ultrasound assessment is used to complement the clinical evaluation, assessing the technical success of treatment, which may determine the long-term evolution of the treated leg. DUS is also ideal for quality control of the initial intervention. Ideally, it is performed by an independent observer. DUS data are objective and reproducible, particularly when obtained and reported using standard methodology, which facilitates reliable collection of serial data in individual patients and in patient groups.

Using a single grading system for clinical outcome that combines symptom scores, clinical findings and DUS findings to define the global outcome will result in an incoherent system, as the relative importance of individual variables may differ. It is therefore appropriate to report separately on clinical outcome parameters and DUS findings in all patients included in prospective studies, focussing on the outcome of treatment of varicose veins, and the effect on other grades of chronic venous disease.

Acknowledgements

The authors like to thank Rebecca Winterborn, Felizitas Pannier and Christina Jeanneret for their valuable collaboration. The authors also thank Ron Slagter for making the diagram (online version).

List of experts who were invited to review this document via Internet: F. Breu (Germany), J. Duque Botero (Colombia), D. Creton (France), B. Eklöf (Sweden), D. Heim (Switzerland), N. Labropoulos (USA), M. Malouf (Australia), N. Morrison (USA), M. Neumann (Netherlands), M. Perrin

(France), N. Pizano (Colombia), E. Rabe (Germany), R. Simkin (Argentina), M. Vandendriessche (Belgium) and C. Wittens (The Netherlands)

Conflict of Interest/Funding

None.

Appendix Supplementary data

Supplementary data related to this article can be found online at [doi:10.1016/j.ejvs.2011.03.013](https://doi.org/10.1016/j.ejvs.2011.03.013).

References

- 1 Redwood NF, Lambert D. Patterns of reflux in recurrent varicose veins. Part 1: evaluation using duplex scanning. *Br J Surg* 1994; **81**:1450–1.
- 2 Bradbury AW, Stonebridge PA, Callam MJ, Walker AJ, Allan PL, Beggs I, et al. Recurrent varicose veins: assessment of the saphenofemoral junction. *Br J Surg* 1994; **81**:373–5.
- 3 Khaira AS, Parnell A. Patterns of reflux in recurrent varicose veins assessed by duplex scanning. *Br J Surg* 1995; **82**:564.
- 4 Labropoulos N, Touloupakis E, Giannoukos AD, Leon M, Katsamouris AD, Nicolaides AN. Recurrent varicose veins: investigation of the pattern and extent of reflux with color flow duplex scanning. *Surgery* 1996; **119**:406–9.
- 5 Wong JK, Duncan JL, Nichols DM. Whole-leg duplex mapping for varicose veins: observations on patterns of reflux in recurrent and primary legs, with clinical correlation. *Eur J Vasc Endovasc Surg* 2003; **25**:267–75.
- 6 van Rij AM, Jiang P, Solomon C, Christie RA, Hill GB. Recurrence after varicose vein surgery: a prospective long-term clinical study with duplex ultrasound scanning and air plethysmography. *J Vasc Surg* 2003; **38**:935–43.
- 7 Jones L, Braithwaite BD, Selwyn D, Cooke S, Earnshaw JJ. Neovascularisation is the principle cause of varicose vein recurrence: results of a randomised trial of stripping the long saphenous vein. *Eur J Vasc Endovasc Surg* 2006; **12**:442–5.
- 8 Coleridge-Smith P, Labropoulos N, Partsch H, Myers K, Nicolaides A, Cavezzi A. Duplex ultrasound investigation of the veins in chronic venous disease of the lower limbs – UIP consensus document. Part I. Basic principles. *Eur J Vasc Endovasc Surg* 2006; **31**:83–92.
- 9 Cavezzi A, Labropoulos N, Partsch H, Ricci S, Caggiati A, Myers K, et al. Duplex ultrasound investigation of the veins in chronic venous disease of the lower limbs – UIP consensus document. Part II. Anatomy. *Eur J Vasc Endovasc Surg* 2006; **31**:288–99.
- 10 De Maeseneer MG, Vandenbroeck CP, Hendriks JM, Lauwers PR, Van Schil PE. Accuracy of duplex evaluation one year after varicose vein surgery to predict recurrence at the saphenofemoral junction after five years. *Eur J Vasc Endovasc Surg* 2005; **29**:308–12.
- 11 Kalra M, Głowiczki P. Fifteen years ago laser was supposed to open arteries, now it is supposed to close veins: what is the reality behind the tool? *Perspect Vasc Surg Endovasc Ther* 2006; **18**:3–8.
- 12 Kundu S, Lurie F, Millward SF, Padberg F, Vedantham S, Elias S, et al. Recommended reporting standards for endovenous ablation for the treatment of venous insufficiency: joint statement of The American Venous Forum and The Society of Interventional Radiology. *J Vasc Surg* 2007; **46**:582–9.

- 13 Breu FX, Guggenbichler S, Wollmann JC. Duplex ultrasound and efficacy criteria in foam sclerotherapy from the 2nd European consensus meeting on foam sclerotherapy 2006 Tegernsee, Germany. *VASA* 2008;**37**:90–5. supplement in *VASA* 2008; **37**(S/71): 3–29.
- 14 Garratt AM, Macdonald LM, Ruta DA, Russell IT, Buckingham JK, Krukowski ZH. Towards measurement of outcome for patients with varicose veins. *Qual Health Care* 1993;**2**:5–10.
- 15 Smith JJ, Garratt AM, Guest M, Greenhalgh RM, Davies AH. Evaluating and improving health-related quality of life in patients with varicose veins. *J Vasc Surg* 1999;**30**:710–9.
- 16 Rutherford RB, Padberg FT, Comerota AJ, Kistner RL, Meissner MH, Moneta GL. Venous severity scoring: an adjunct to venous outcome assessment. *J Vasc Surg* 2000;**31**:1307–12.
- 17 Eklöf B, Rutherford RB, Bergan JJ, Carpentier PH, Gloviczki P, Kistner RL, et al. American venous forum international ad hoc committee for revision of the CEAP classification. *J Vasc Surg* 2004;**40**:1248–52.
- 18 Guex JJ, Zimmet SE, Boussetta S, Nguyen C, Taieb C. Construction and validation of a patient-reported outcome dedicated to chronic venous disorders: SQOR-V (specific quality of life and outcome response – venous). *J Mal Vasc* 2007;**32**:135–47.
- 19 Vasquez MA, Munschauer CE. Venous clinical severity score and quality-of-life assessment tools: application to vein practice. *Phlebology* 2008;**23**:259–75.
- 20 Perrin MR, Guex JJ, Ruckley CV, de Palma RG, Royle JP, Eklof B, et al. The REVAS group. Recurrent varicose veins after surgery (REVAS), a consensus document. *Cardiovasc Surg* 2000;**8**:233–45.
- 21 Perrin MR, Labropoulos N, Leon LR. Presentation of the patient with recurrent varices after surgery (REVAS). *J Vasc Surg* 2006; **43**:327–34.
- 22 Eklöf B, Perrin M, Delis KT, Rutherford RB, Gloviczki P. Updated terminology of chronic venous disorders: the VEIN-TERM transatlantic interdisciplinary document. *J Vasc Surg* 2009;**49**: 498–501.
- 23 Labropoulos N, Kokkosis AA, Spentzouris G, Gasparis AP, Tassiopoulos AK. The distribution and significance of varicosities in the saphenous trunks. *J Vasc Surg* 2010;**51**:96–103.
- 24 Neglén P, Egger JF, Olivier J, Raju S. Hemodynamic and clinical impact of ultrasound derived venous reflux parameters. *J Vasc Surg* 2004;**40**:303–10.
- 25 Jeanneret C, Labs KH, Aschwanden M, Bollinger A, Hoffmann U, Jäger K. Physiologic reflux and venous diameter change in the proximal lower limb during a standardised Valsalva manoeuvre. *Eur J Vasc Endovasc Surg* 1999;**17**:398–403.
- 26 Yamaki T, Nozaki M, Sakurai H, Takeuchi M, Soejima K, Kono T. Comparison of manual compression release with distal pneumatic cuff maneuver in the ultrasonic evaluation of superficial venous insufficiency. *Eur J Vasc Endovasc Surg* 2006;**32**:462–7.
- 27 Lemasle P, Uhl JF, Lefebvre-Vilardebo M, Baud JM, Gillot C. Veines lympho-ganglionnaires inguinales. Aspects anatomiques et échographiques. Conséquences sur la définition de la néogenèse. Conséquences thérapeutiques. *Phlébologie* 1999;**52**:263–9.
- 28 Georgiev M. The preoperative duplex examination. *Dermatol Surg* 1998;**24**:433–40.
- 29 van Rij AM, Chai J, Hill GB, Christie RA. Incidence of deep vein thrombosis after varicose vein surgery. *Br J Surg* 2004;**91**: 1582–5.
- 30 Puggioni A, Kalra M, Carmo M, Mozes G, Gloviczki P. Endovenous laser therapy and radiofrequency ablation of the great saphenous vein: analysis of early efficacy and complications. *J Vasc Surg* 2005;**42**:488–93.
- 31 Mozes G, Kalra M, Carmo M, Swenson L, Gloviczki P. Extension of saphenous thrombus into the femoral vein: a potential complication of new endovenous ablation techniques. *J Vasc Surg* 2005;**41**:130–5.
- 32 Knipp BS, Blackburn SA, Bloom JR, Fellows E, Laforge W, Pfeifer JR, et al Michigan Venous Study Group. Endovenous laser ablation: venous outcomes and thrombotic complications are independent of the presence of deep venous insufficiency. *Vasc Surg* 2008;**48**:1538–45.
- 33 Myers KA, Jolley D. Factors affecting the risk of deep venous occlusion after ultrasound-guided sclerotherapy for varicose veins. *Eur J Vasc Endovasc Surg* 2008;**36**:602–5.
- 34 Pares JO, Juan J, Tellez R, Mata A, Moreno C, Quer FX, et al. Varicose vein surgery: stripping versus the CHIVA method. A randomized controlled trial. *Ann Surg* 2010;**251**:624–31.
- 35 Min RJ, Khilnani N, Zimmet S. Endovenous laser treatment of saphenous vein reflux: long term results. *J Vasc Interv Radiol* 2003;**14**:991–6.
- 36 Merchant RF, Pichot O Closure Study Group. Long-term outcomes of endovenous radiofrequency obliteration of saphenous reflux as a treatment for superficial venous insufficiency. *J Vasc Surg* 2005;**42**:502–9.
- 37 De Maeseneer MG, Ongena KP, Van den Brande F, Van Schil PE, De Hert SG, Eyskens EJ. Duplex ultrasound assessment of neo-vascularization after sapheno-femoral or sapheno-popliteal junction ligation. *Phlebology* 1997;**12**:64–8.
- 38 Earnshaw JJ, Davies B, Harradine K, Heather BP. Preliminary results of PTFE patch saphenoplasty to prevent neo-vascularization leading to recurrent varicose veins. *Phlebology* 1998;**13**:10–3.
- 39 De Maeseneer MG, Giuliani DR, Van Schil PE, De Hert SG. Can interposition of a silicone implant after sapheno-femoral ligation prevent recurrent varicose veins? *Eur J Vasc Endovasc Surg* 2002;**24**:445–9.
- 40 De Maeseneer MG, Philipsen TE, Vandenbroeck CP, Lauwers PR, Hendriks JM, De Hert SG, et al. Closure of the cribriform fascia: an efficient anatomical barrier against postoperative neo-vascularisation at the saphenofemoral junction? A prospective study. *Eur J Vasc Endovasc Surg* 2007;**34**:361–6.
- 41 van Rij AM, Jones GT, Hill BG, Amer M, Thomson IA, Pettigrew RA, et al. Mechanical inhibition of angiogenesis at the saphenofemoral junction in the surgical treatment of varicose veins. Early results of a blinded randomized controlled trial. *Circulation* 2008;**118**:66–74.
- 42 Agus GB, Magi GIEWG. The first 1000 cases of Italian Endovenous-laser Working Group (IEGW). Rationale, and long-term outcomes for the 1999–2003 period. *Int Angiol* 2006;**25**:209–15.
- 43 Myers KA, Jolley D, Clough A, Kirwan J. Outcome of ultrasound-guided sclerotherapy for varicose veins: medium-term results assessed by ultrasound surveillance. *Eur J Vasc Endovasc Surg* 2007;**33**:116–21.
- 44 Dwerryhouse S, Davies B, Harradine K, Earnshaw JJ. Stripping the long saphenous vein reduces the rate of reoperation for recurrent varicose veins. *J Vasc Surg* 1999;**29**:589–92.
- 45 Creton D. Surgery for recurrent sapheno-femoral incompetence using expanded polytetrafluoroethylene patch interposition in front of the femoral vein: long-term outcome in 119 extremities. *Phlebology* 2002;**16**:93–7.
- 46 Winterborn RJ, Foy C, Earnshaw JJ. Causes of varicose vein recurrence: late results of a randomised controlled trial of stripping the long saphenous vein. *J Vasc Surg* 2004;**40**:634–9.
- 47 De Maeseneer MG, Vandenbroeck CP, Van Schil PE. Silicone patch saphenoplasty to prevent repeat recurrence after surgery to treat recurrent saphenofemoral incompetence: long-term follow-up study. *J Vasc Surg* 2004;**40**:98–105.
- 48 Lane RJ, Graiche JA, Coroneos JC, Cuzzilla ML. Long-term comparison of external valvular stenting and stripping of varicose veins. *Aust NZ J Surg* 2003;**73**:605–9.
- 49 Hartmann K, Klode J, Pfister R, Toussaint M, Weingart I, Waldermann F, et al. Recurrent varicose veins: sonography-based re-examination of 210 patients 14 years after ligation and saphenous vein stripping. *VASA* 2006;**35**:21–6.
- 50 Carandina S, Mari C, De Palma M, Marcellino MG, Cisno C, Legnaro A, et al. Varicose vein stripping vs. haemodynamic

- correction (CHIVA): a long term randomised trial. *Eur J Vasc Endovasc Surg* 2008;**35**:230–7.
- 51 Fischer R, Linde N, Duff C, Jeanneret C, Chandler JG, Seeber P. Late recurrent saphenofemoral junction reflux after ligation and stripping of the greater saphenous vein. *J Vasc Surg* 2001;**34**:236–40.
- 52 Fischer R, Chandler JG, De Maeseneer MG, Frings N, Lefebvre-Vilardebo M, Earnshaw JJ, et al. The unresolved problem of recurrent saphenofemoral reflux. *J Am Coll Surg* 2002;**195**: 80–94.
- 53 van Rij AM, Jones GT, Hill GB, Jiang P. Neovascularization and recurrent varicose veins: more histologic and ultrasound evidence. *J Vasc Surg* 2004;**40**:296–302.
- 54 Stücker M, Netz K, Breuckmann F, Altmeyer P, Mumme A. Histomorphologic classification of recurrent saphenofemoral reflux. *J Vasc Surg* 2004;**39**:816–22.
- 55 De Maeseneer MGR. The role of postoperative neovascularisation in recurrence of varicose veins. *Acta Chir Belg* 2004;**104**: 281–7.
- 56 Egan B, Donnelly M, Bresnihan M, Tierney S, Feeley M. Neovascularization: an “innocent bystander” in recurrent varicose veins. *J Vasc Surg* 2006;**44**:1279–84.
- 57 Geier B, Mumme A, Hummel T, Marpe B, Stücker M, Ascitto G. Validity of duplex-ultrasound in identifying the cause of groin recurrence after varicose vein surgery. *J Vasc Surg* 2009;**49**: 968–72.
- 58 Lemasle Ph, Lefebvre-Vilardebo M, Uhl JF, Vin F, Baud JM. Récidive variqueuse postopératoire: et si la néo-vascularisation inguinale post-chirurgicale n’était que le développement d’un réseau pré-existant? *Phlébologie* 2009;**62**:42–8.
- 59 Pittaluga P, Chastanet S, Guex JJ. Great saphenous vein stripping with preservation of sapheno-femoral confluence: hemodynamic and clinical results. *J Vasc Surg* 2008;**47**:1300–4.
- 60 Casoni P. Is still the crossectomy the first obliged step in varicose vein surgery? 6 yrs follow up in a randomized study in 124 legs without inguinal dissection. *Int Angiol* 2009;**28**(Suppl. 1): 112–3.
- 61 Munasinghe A, Smith C, Klanifard B, Price BA, Holdstock JM, Whiteley MS. Strip-track revascularization after stripping of the great saphenous vein. *Br J Surg* 2007;**94**:840–3.
- 62 De Maeseneer MG, De Hert SG, Van Schil PE, Vanmaele RG, Eyskens EJ. Preoperative colour coded duplex examination of the saphenopopliteal junction in recurrent varicosis of the short saphenous vein. *Cardiovasc Surg* 1993;**1**:686–9.
- 63 Tong Y, Royle J. Recurrent varicose veins after short saphenous vein surgery: a duplex ultrasound study. *Cardiovasc Surg* 1996;**4**:364–7.
- 64 Perrin M, Gillet JL. Management of recurrent varices at the popliteal fossa after surgical treatment. *Phlebology* 2008;**23**: 64–8.
- 65 O’Hare JL, Vandenbroeck CP, Whitman B, Campbell B, Heather BP, Earnshaw JJ. Joint Vascular Research Group. *J Vasc Surg* 2008;**48**:669–73.
- 66 Cavezzi A, Tarabini C, Collura M, Sigismondi G, Barboni MG, Carigi V. Hémodynamique de la jonction saphéno-poplitée: évaluation par écho-doppler couleur. *Phlébologie* 2002;**55**: 309–16.
- 67 van Rij AM, Hill G, Gray C, Christie R, Macfarlane J, Thomson I. A prospective study of the fate of venous leg perforators after varicose vein surgery. *J Vasc Surg* 2005;**42**:1156–62.
- 68 García-Gimeno M, Rodríguez-Camarero S, Tagarro-Villalba S, Ramalle-Gomara E, González-González E, Vaquero Puerta C. Duplex mapping of 2036 primary varicose veins. *J Vasc Surg* 2009;**49**:681–9.
- 69 Stuart WP, Adam DJ, Allan PL, Ruckley C, Bradbury AW. Saphenous surgery does not correct perforator incompetence in the presence of deep venous reflux. *J Vasc Surg* 1998;**28**: 834–8.
- 70 Recek C. Impact of calf perforators on the venous hemodynamics in primary varicose veins. *J Cardiovasc Surg (Torino)* 2006;**47**:629–35.
- 71 Lurie F, Creton D, Eklof B, Kabnick LS, Kistner RL, Pichot O, et al. Prospective randomised study of endovenous radiofrequency ablation (closure procedure) versus ligation and stripping in a selected patient population (EVOLVE Study). *J Vasc Surg* 2003;**38**:207–14.
- 72 Pichot O, Sessa C, Chandler JG, Nuta M, Perrin M. Role of duplex imaging in endovenous obliteration for primary venous insufficiency. *Endovasc Ther* 2000;**7**:451–9.
- 73 Proebstle TM, Vago B, Alm J, Göckeritz O, Lebard C, Pichot O, for the Closure Fast Clinical Study Group. Treatment of the incompetent great saphenous vein by endovenous radiofrequency powered segmental thermal ablation: first clinical experience. *J Vasc Surg* 2008;**47**:151–6.
- 74 Puggioni A, Marks N, Hingorani A, Shiferson A, Alhalbouni S, Ascher E. The safety of radiofrequency ablation of the great saphenous vein in patients with previous venous thrombosis. *J Vasc Surg* 2009;**49**:1248–55.
- 75 Sharif MA, Soong CV, Lau LL, Lee B, Hannon RJ. Endovenous laser treatment for long saphenous vein incompetence. *Br J Surg* 2006;**93**:831–5.
- 76 Rasmussen LH, Bjoern L, Lawaetz M, Lawaetz B, Blemings A, Eklöf B. Randomised clinical trial comparing endovenous laser ablation with stripping of the great saphenous vein: clinical outcome and recurrence after 2 years. *Eur J Vasc Endovasc Surg* 2010;**39**:630–5.
- 77 Theivacumar NS, Dellagrammaticas D, Beale RJ, Mavor AI, Gough MJ. Factors influencing the effectiveness of endovenous laser ablation (EVLA) in the treatment of great saphenous vein reflux. *Eur J Vasc Endovasc Surg* 2008;**35**:119–23.
- 78 Proebstle TM, Gul D, Kargl A, Knop J. Endovenous laser treatment of the lesser saphenous vein with a 940-nm diode laser: early results. *Dermatol Surg* 2003;**29**:357–61.
- 79 Park SW, Hwang JJ, Yun IJ, Lee SA, Kim JS, Chang SH, et al. Endovenous laser ablation of the incompetent small saphenous vein with a 980-nm diode laser: our experience with 3years follow-up. *Eur J Vasc Endovasc Surg* 2008;**36**:738–42.
- 80 Desmyttere J, Grard C, Stalnickiewicz G, Wassmer B, Mordon S. Endovenous laser ablation (980 nm) of the small saphenous vein in a series of 147 limbs with a 3-year follow-up. *Eur J Vasc Endovasc Surg* 2010;**39**:99–103.
- 81 Coleridge Smith P. Chronic venous disease treated by ultrasound guided foam sclerotherapy. *Eur J Vasc Endovasc Surg* 2006;**32**: 577–83.
- 82 Hamel-Desnos C, Ouvry P, Benigni JP, Boitelle G, Schadeck M, Desnos P, et al. Comparison of 1% and 3% polidocanol foam in ultrasound guided sclerotherapy of the great saphenous vein: a randomised double-blind trial with 2 year-follow-up. “The 3/1 study”. *Eur J Vasc Endovasc Surg* 2007;**34**:723–9.
- 83 Parsi K. Catheter-directed sclerotherapy. *Phlebology* 2009;**24**: 98–107.
- 84 Hingorani A, Ascher E, Markevich N, Schutzer RW, Kallakuri S, Hou A, et al. Deep venous thrombosis after radiofrequency ablation of greater saphenous vein: a word of caution. *J Vasc Surg* 2004;**40**:500–4.
- 85 Marsh P, Price BA, Holdstock J, Harrison C, Whiteley MS. Deep vein thrombosis (DVT) after venous thermoablation techniques: rate of endovenous heat-induced thrombosis (EHIT) and classical DVT after radiofrequency and endovenous laser ablation in a single centre. *Eur J Vasc Endovasc Surg* 2010;**40**: 521–7.
- 86 Lawrence PF, Chandra A, Wu M, Rigberg D, De Rubertis D, Gelabert H, et al. Classification of proximal endovenous closure levels and treatment algorithm. *J Vasc Surg* 2010;**52**:388–93.
- 87 Franceschi C, Zamboni P. *Principles of venous haemodynamics*. New York: Nova Science Publisher; 2009.

- 88 Theivacumar NS, Darwood R, Gough MJ. Neovascularisation and recurrence after varicose vein treatment for sapheno-femoral and great saphenous vein reflux: a comparison of surgery and endovenous laser ablation. *Eur J Vasc Endovasc Surg* 2009;**38**: 203–7.
- 89 Pichot O, Kabnick L, Creton D, Merchant RF, Schuller-Petrovic S, Chandler JG. Duplex ultrasound scan findings two years after great saphenous vein radiofrequency endovenous obliteration. *J Vasc Surg* 2004;**39**:189–95.
- 90 Weiss RA. Comparison of endovenous radiofrequency versus 810 nm diode laser occlusion of large veins in an animal model. *Dermatol Surg* 2002;**28**:56–61.
- 91 Schmedt CG, Meissner OA, Hunger K, Babaryka G, Ruppert V, Sadeghi-Azandaryani M, et al. Evaluation of endovenous radiofrequency ablation and laser therapy with endoluminal optical coherence tomography in an ex vivo model. *J Vasc Surg* 2007;**45**:1047–58.
- 92 Theivacumar NS, Dellagrammaticas D, Mavor AI, Gough MJ. Endovenous laser ablation: does standard above-knee great saphenous vein ablation provide optimum results in patients with both above- and below-knee reflux? A randomized controlled trial. *J Vasc Surg* 2008;**48**:173–8.
- 93 Peden E, Lumsden A. Radiofrequency ablation of incompetent perforating veins. *Perspect Vasc Surg Endovasc Ther* 2007;**19**: 73–7.
- 94 Elias S, Peden E. Ultrasound-guided percutaneous ablation for the treatment of perforating vein incompetence. *Vascular* 2007;**15**:281–9.